arXiv:1301.2361v1 [math.GT] 11 Jan 2013

LEFT-ORDERABLE FUNDAMENTAL GROUP AND DEHN SURGERY ON GENUS ONE TWO-BRIDGE KNOTS

RYOTO HAKAMATA AND MASAKAZU TERAGAITO

Abstract. For any hyperbolic genus one 2-bridge knot in the 3-sphere, we show that the resulting manifold by r-surgery on the knot has left-orderable fundamental group if the slope r lies in some range which depends on the knot.

1. Introduction

A non-trivial group G is said to be *left-orderable* if it admits a strict total ordering which is invariant under left-multiplication. Thus, if $g < h$ then $fg < fh$ for any $f, g, h \in G$. Many fundamental groups of 3-manifolds are known to be left-orderable. For examples, all knot and link groups are left-orderable. Boyer, Gordon and Watson [3] propose a conjecture that an irreducible rational homology 3-sphere Y is an L-space if and only if $\pi_1 Y$ is not left-orderable. An L-space is a rational homology 3-sphere whose Heegaard–Floer homology $\widehat{HF}(Y)$ has rank $|H_1(Y;\mathbb{Z})|$ ([16]). Recently, L-spaces become the object of interest, and it is an open problem to characterize L-spaces without mentioning Heegaard–Floer homology. The affirmative answer to the above conjecture will give an algebraic characterization of L-spaces.

On the other hand, irreducible rational homology spheres are obtained by Dehn surgery on knots in the 3-sphere S^3, in plenty. For a knot K, we call a slope r *left-orderable* if the resulting manifold $K(r)$ by r-surgery on K has left-orderable fundamental group. In this paper, a slope is sometimes identified with its parameter in $\mathbb{Q} \cup \{1/0\}$. In particular, the meridional slope corresponds to $1/0$. It is known that any hyperbolic 2-bridge knot does not admit Dehn surgery yielding an L-space ([16]). Hence any slope but $1/0$ is expected to be left-orderable for a hyperbolic 2-bridge knot, if we support the above conjecture. In this direction, Boyer, Gordon and Watson [3] proved that for the figure-eight knot, if $r \in (-4, 4)$, then r is left-orderable. Later, Clay, Lidman and Watson [6] showed $r = \pm 4$ are also left-orderable. These results were extended to all hyperbolic twist knots [10, 11, 19]. Also, Tran [20] further extended the range of left-orderable slopes for twist knots.

2010 *Mathematics Subject Classification.* Primary 57M25; Secondary 06F15.

Key words and phrases. left-ordering, Dehn surgery, two-bridge knot.

The second author is partially supported by Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research (C), 22540088.

The purpose of this paper is to give new ranges of left-orderable slopes for all hyperbolic genus one, 2-bridge knots. As a by-product, we obtain a wider range of left-orderable slopes for (positive) twist knots than that in [11].

For non-zero integers m and n, let $K(m,n)$ be the 2-bridge knot $S(4mn+1,2m)$ in Schubert's normal form as illustrated in Figure 1. In Figure 1, the twists in the vertical box are left-handed (resp. right-handed) if $m>0$ (resp. $m<0$), but those in the horizontal box are right-handed (resp. left-handed) if $n>0$ (resp. $n<0$). Thus $K(1,-1)$ is the trefoil, and $K(1,1)$ is the figure-eight knot. By symmetry, $K(m,n)$ and $K(-n,-m)$ are isotopic. Except the trefoil, $K(m,n)$ is hyperbolic. It is also well known that any genus one 2-bridge knot is equivalent to $K(m,n)$ for some m,n (see [5]).



FIGURE 1. The knot $K(m,n)$

Theorem 1.1. *Let $K=K(m,n)$ be a hyperbolic genus one 2-bridge knot $S(4mn+1,2m)$ in the 3-sphere S^3 as illustrated in Figure 1. Let I be the interval defined by*

$$I=\begin{cases}(-4n,4m) & \text{if } m>0 \text{ and } n>0,\\ (4m,-4n) & \text{if } m<0 \text{ and } n<0,\\ [0,\max\{4m,-4n\}) & \text{if } m>0 \text{ and } n<0,\\ (\min\{4m,-4n\},0] & \text{if } m<0 \text{ and } n>0.\end{cases}$$

Then any slope in I is left-orderable. That is, $\pi_1(K(r))$ is left-orderable.

We remark that $K(1,-1)$ and $K(-1,1)$ are trefoils, so they are excluded in the statement here.

In a previous paper [11], we showed that any hyperbolic twist knot $K(1,n)$ ($n\neq -1$) admits a range $[0,4]$ of left-orderable slopes. Theorem 1.1 gives a partial improvement of this result.

Corollary 1.2. *Let $K=K(1,n)$ be a hyperbolic n-twist knot for $n>0$. If $I=(-4n,4]$, then any slope in I is left-orderable.*

This range of Corollary 1.2 coincides with that in [20]. Also, Tran informed us that he obtained a similar result to Theorem 1.1. We would like to thank Anh T. Tran for informing his result.

2. Knot groups and Two sequences of polynomials

Let $K = K(m,n)$ and let $G = \pi_1(S^3 - K)$ be its knot group. We always assume that $m \neq 0$ and $n \neq 0$, unless specified otherwise.

Proposition 2.1. *The knot group G admits a presentation*

$$G = \langle x, y \mid w^n x = y w^n \rangle,$$

where x and y are meridians and $w = (xy^{-1})^m (x^{-1}y)^m$. Furthermore, the longitude $\mathcal{L}$ is given as $\mathcal{L} = w_*^n w^n$, where $w_* = (yx^{-1})^m (y^{-1}x)^m$ is obtained from w by reversing the order of letters.

This is slightly different from that in [13, Proposition 1], but both are isomorphic.

Proof. We use a surgery diagram of K as illustrated in Figure 2, where $1/m$–surgery and $-1/n$–surgery are performed along the second and third components, respectively. Let μ_i and λ_i be the meridian and longitude of the ith component.



Figure 2. A surgery diagram of K

First, $y = \mu_3^{-1} x \mu_3$, $z = \mu_2^{-1} y \mu_2$, $\lambda_2 = x^{-1}y$ and $\lambda_3 = yz^{-1}$. By $1/m$–surgery on the second component yields a relation $\lambda_2^m \mu_2 = 1$, so $\mu_2 = \lambda_2^{-m}$. Similarly, $-1/n$–surgery on the third component gives $\mu_3 = \lambda_3^n$. Thus

$$\begin{aligned} \lambda_3 &= y\mu_2^{-1}y^{-1}\mu_2 = y\lambda_2^m y^{-1}\lambda_2^{-m} \\ &= y(x^{-1}y)^m y^{-1}(y^{-1}x)^m = (yx^{-1})^m(y^{-1}x)^m. \end{aligned}$$

Then the relation $y = \mu_3^{-1} x \mu_3$ gives

$$y[(x^{-1}y)^m(xy^{-1})^m]^n = [(x^{-1}y)^m(xy^{-1})^m]^n y. \tag{2.1}$$

Set $w = (xy^{-1})^m(x^{-1}y)^m$. Since $[(x^{-1}y)^m(xy^{-1})^m]^n = (x^{-1}y)^m w^n (x^{-1}y)^{-m}$, (2.1) changes to $y(x^{-1}y)^m w^n (x^{-1}y)^{-m} = (x^{-1}y)^m w^n (x^{-1}y)^{-m} x$. By $y(x^{-1}y)^m =$

$(yx^{-1})^m y$, we have $(yx^{-1})^m yw^n(x^{-1}y)^{-m} = (x^{-1}y)^m w^n (x^{-1}y)^{-m}x$. Thus,

$$\begin{aligned} yw^n &= (xy^{-1})^m(x^{-1}y)^m w^n(x^{-1}y)^{-m}x(x^{-1}y)^m \\ &= w^{n+1}(x^{-1}y)^{-m}x(x^{-1}y)^m \\ &= w^n(xy^{-1})^m(x^{-1}y)^m(x^{-1}y)^{-m}x(x^{-1}y)^m \\ &= w^n(xy^{-1})^m x(x^{-1}y)^m \\ &= w^n x. \end{aligned}$$

Set $w_* = (yx^{-1})^m(y^{-1}x)^m$. Then $\lambda_3 = w_*$. The longitude is $\mathcal{L} = \mu_3\mu_2\mu_3^{-1}\mu_2^{-1} = w_*^n\mu_2 w_*^{-n}\mu_2^{-1}$. We have

$$\begin{aligned} \mu_2 w_*^{-n}\mu_2^{-1} &= (x^{-1}y)^{-m}[(yx^{-1})^m(y^{-1}x)^m]^{-n}(x^{-1}y)^m \\ &= (x^{-1}y)^{-m}[(x^{-1}y)^m(xy^{-1})^m]^n(x^{-1}y)^m \\ &= [(xy^{-1})^m(x^{-1}y)^m]^n \\ &= w^n. \end{aligned}$$

Thus $\mathcal{L} = w_*^n w^n$. □

To describe the Riley polynomial of K in Section 3, we prepare two sequences of polynomials with a single variable s.

For non-negative integer m, let $f_m \in \mathbb{Z}[s]$ be defined by the recursion

$$f_{m+2} - (s+2)f_{m+1} + f_m = 0 \tag{2.2}$$

with initial conditions $f_0 = 1$ and $f_1 = s+1$. Also, let $g_m \in \mathbb{Z}[s]$ be defined by the same recursion

$$g_{m+2} - (s+2)g_{m+1} + g_m = 0 \tag{2.3}$$

with slightly different initial conditions $g_0 = 1$ and $g_1 = s+2$. We remark that g_m is equivalent to the Chebyshev polynomial of the second kind.

Lemma 2.2. *The closed formulas for f_m and g_m are*

$$f_m = \sum_{i=0}^{m}\binom{m+i}{m-i}s^i, \quad g_m = \sum_{i=0}^{m}\binom{m+1+i}{m-i}s^i.$$

In particular, all coefficients of f_m and g_m are positive integers, and the degree of f_m and g_m is m. Also, f_m and g_m are monic.

Proof. These easily follow from the inductive argument by using the recursive formulas. □

Set $f_{-m} = f_{m-1}$ for $m \geq 1$, and set $g_{-1} = 0$ and $g_{-m} = -g_{m-2}$ for $m \geq 2$. Thus we have defined f_m and g_m for any integer m. In particular, the recursions (2.2) and (2.3) hold for all integers.

Lemma 2.3. *For any integer m, the polynomials f_m, g_m satisfy the following relations.*

1. $f_m + g_{m-1} = g_m$.
2. $f_m + sg_m = f_{m+1}$.
3. $f_m^2 = sg_m g_{m-1} + 1$.

Proof. These are easily proved by induction. We prove (3) only here. Clearly, it holds when $m = 0$. Assume $f_m^2 = sg_m g_{m-1} + 1$. From (1) and (2),

$$\begin{aligned} f_{m+1}^2 &= (f_m + sg_m)^2 \\ &= f_m^2 + 2sf_m g_m + s^2 g_m^2 \\ &= (sg_m g_{m-1} + 1) + 2sf_m g_m + s^2 g_m^2 \\ &= (sg_m g_{m-1} + 1) + 2s(g_m - g_{m-1})g_m + s^2 g_m^2 \\ &= (s^2 + 2s)g_m^2 - sg_m g_{m-1} + 1 \\ &= s((s+2)g_m - g_{m-1})g_m + 1 \\ &= sg_{m+1}g_m + 1. \end{aligned}$$

Similarly, $f_{m-1}^2 = (f_m - sg_{m-1})^2 = sg_{m-1}g_{m-2} + 1$. □

Lemma 2.4. *If a positive real number is substituted to s, then we have the following inequalities.*

1. *For $m > 0$, $sg_{m-1} < 4f_m$.*
2. *For $m < 0$, $-sg_{m-1} < 4f_{m-1}$.*
3. *For $m < 0$, $-2sg_{m-1} < 3f_{m-1}$.*

Proof. (1) Since $sg_{m-1} = f_m - f_{m-1}$ by Lemma 2.3, this follows from $3f_m + f_{m-1} > 0$. (2) is equivalent to (1). (3) From $sg_{m-1} = f_m - f_{m-1}$, it follows from $2f_m + f_{m-1} > 0$. □

3. Riley polynomials

In this section, we calculate the Riley polynomial of K. Let s and t be real numbers such that $s > 0$ and $t > 1$. Let $\rho : G \to SL_2(\mathbb{R})$ be a representation of G defined by

$$\rho(x) = \begin{pmatrix} \sqrt{t} & 1/\sqrt{t} \\ 0 & 1/\sqrt{t} \end{pmatrix}, \quad \rho(y) = \begin{pmatrix} \sqrt{t} & 0 \\ -s\sqrt{t} & 1/\sqrt{t} \end{pmatrix}.$$

By [17], ρ gives a non-abelian representation if s and t are a pair of solutions of the Riley polynomial. Recall that $w = (xy^{-1})^m(x^{-1}y)^m$. Let $W = \rho(w)$ and $z_{i,j}$ be the (i,j)-entry of W^n. Then the Riley polynomial of K is given by $\phi_K(s,t) = z_{1,1} + (1-t)z_{1,2}$. (See also [8].)

Since s and t are limited to be positive real numbers in our setting, it is not obvious that there exist solutions for Riley's equation $\phi_K(s,t) = 0$. However, this will be verified in Proposition 4.2 under some condition. In fact, we can choose t so that $t > 1$ and $s + 2 < t + 1/t < s + 2 + 4/(sg_{m-1}^2)$ for any $s > 0$. We temporarily assume that s and t are chosen to satisfy $\phi_K(s,t) = 0$. See Example 3.7 when $n = \pm 1$.

Proposition 3.1. *For* $W = \rho(w)$, *we have*

$$W = \begin{pmatrix} f_m^2 - stg_{m-1}^2 & f_{m-1}g_{m-1} - \frac{f_m g_{m-1}}{t} \\ sf_m g_{m-1} - stf_{m-1}g_{m-1} & f_{m-1}^2 - \frac{s}{t}g_{m-1}^2 \end{pmatrix}.$$

Proof. We prove by induction on m. (For the knot $K(m,n)$, we assume $m \neq 0$. However, this proposition holds even for $m = 0$.)

If $m = 0$, then $w = 1$, so W is the identity matrix. It is easy to check that the claim holds.

Assume the conclusion for m. Note

$$\rho(xy^{-1}) = \begin{pmatrix} s+1 & 1 \\ s & 1 \end{pmatrix}, \ \rho(x^{-1}y) = \begin{pmatrix} s+1 & -1/t \\ -st & 1 \end{pmatrix}.$$

Calculate the product

$$\begin{pmatrix} s+1 & 1 \\ s & 1 \end{pmatrix} \begin{pmatrix} f_m^2 - stg_{m-1}^2 & f_{m-1}g_{m-1} - \frac{f_m g_{m-1}}{t} \\ sf_m g_{m-1} - stf_{m-1}g_{m-1} & f_{m-1}^2 - \frac{s}{t}g_{m-1}^2 \end{pmatrix} \begin{pmatrix} s+1 & -1/t \\ -st & 1 \end{pmatrix}.$$

By using Lemma 2.3, each entry is identified as desired. For example, the $(1,1)$-entry is

$$\begin{aligned}(f_m + s(f_m + g_{m-1}))^2 - st(f_{m-1} + g_{m-1} + sg_{m-1})^2 &= (f_m + sg_m)^2 - st(f_{m-1} + (s+1)g_{m-1})^2 \\ &= f_{m+1}^2 - st(f_m + g_{m-1})^2 \\ &= f_{m+1}^2 - stg_m^2.\end{aligned}$$

Similarly,

$$\rho(yx^{-1}) = \begin{pmatrix} 1 & -1 \\ -s & s+1 \end{pmatrix}, \ \rho(y^{-1}x) = \begin{pmatrix} 1 & 1/t \\ st & s+1 \end{pmatrix}.$$

Then, calculating the product

$$\begin{pmatrix} 1 & -1 \\ -s & s+1 \end{pmatrix} \begin{pmatrix} f_m^2 - stg_{m-1}^2 & f_{m-1}g_{m-1} - \frac{f_m g_{m-1}}{t} \\ sf_m g_{m-1} - stf_{m-1}g_{m-1} & f_{m-1}^2 - \frac{s}{t}g_{m-1}^2 \end{pmatrix} \begin{pmatrix} 1 & 1/t \\ st & s+1 \end{pmatrix}$$

gives the conclusion again. □

Let $\lambda_\pm \in \mathbb{C}$ be the eigenvalues of W.

Lemma 3.2. $\lambda_+ \neq \lambda_-$.

Proof. Since $\det W = 1$, $\lambda_+\lambda_- = 1$. If $\lambda_+ = \lambda_-$, then $\lambda_+ = \lambda_- = \pm 1$. Then $\mathrm{tr}W = \pm 2$.

On the other hand, $\mathrm{tr}W = f_m^2 + f_{m-1}^2 - s(t+1/t)g_{m-1}^2 = s(s+2-t-1/t)g_{m-1}^2 + 2$ by Proposition 3.1 and Lemma 2.3. Since $g_{m-1} \neq 0$, $s > 0$ and $s + 2 < t + 1/t$,

$\mathrm{tr}W < 2$. If $\mathrm{tr}W = -2$, then $s(t+1/t-s-2)g_{m-1}^2 = 4$, so $t+1/t = s+2+4/(sg_{m-1}^2)$. But this is impossible by our choice of t, remarked before Proposition 3.1. Hence $\lambda_+ \neq \lambda_-$. □

Let $w_{i,j}$ be the (i,j)-entry of W, and let $P = \begin{pmatrix} w_{1,2} & w_{1,2} \\ \lambda_+ - w_{1,1} & \lambda_- - w_{1,1} \end{pmatrix}$.

Lemma 3.3. $w_{1,2} \neq 0$.

Proof. Assume $w_{1,2} = 0$. By Proposition 3.1, $w_{1,2} = f_{m-1}g_{m-1} - f_m g_{m-1}/t$. Since $m \neq 0$, $g_{m-1} \neq 0$. Hence $f_m = tf_{m-1}$.

From the recursion (2.2), $tf_{m-1} - (s+2)f_{m-1} + f_{m-2} = 0$. Since $f_{m-1} \neq 0$, we have $t = s+2-f_{m-2}/f_{m-1}$. Thus $t < s+2$, contradicting our choice of t again. □

The next will be used in Section 5.

Lemma 3.4. $w_{2,1} = -stw_{1,2}$. *Hence* $w_{2,1} \neq 0$.

Proof. This immediately follows from Proposition 3.1 and Lemma 3.3. □

By Lemmas 3.2 and 3.3, $\det P = -w_{1,2}(\lambda_+ - \lambda_-) \neq 0$. A direct calculation shows $P^{-1}WP = \begin{pmatrix} \lambda_+ & 0 \\ 0 & \lambda_- \end{pmatrix}$.

Set $\tau = \mathrm{tr}W$. For any positive integer k, let $\tau_k = \tau^{k-1} + \tau^{k-3} + \cdots + \tau^{3-k} + \tau^{1-k}$. If $\lambda_+ \neq \lambda_-$, then $\tau_k = (\lambda_+^k - \lambda_-^k)/(\lambda_+ - \lambda_-)$. Let $\tau_0 = 0$ and $\tau_{-k} = -\tau_k$. Then a recursion $\tau_{k+1} - \tau\tau_k + \tau_{k-1} = 0$ holds for any integer k.

Lemma 3.5. *For* $W^n = \rho(w^n)$, *we have*

$$W^n = \begin{pmatrix} w_{1,1}\tau_n - \tau_{n-1} & w_{1,2}\tau_n \\ w_{2,1}\tau_n & \tau_{n+1} - w_{1,1}\tau_n \end{pmatrix}.$$

Proof. When $n = \pm 1$, it is easy to see the conclusion. Let $|n| > 1$. This easily follows from $W^n = P\begin{pmatrix} \lambda_+^n & 0 \\ 0 & \lambda_-^n \end{pmatrix}P^{-1}$. □

Proposition 3.6. *The Riley polynomial of* K *is*

$$\phi_K(s,t) = (\tau_{n+1} - \tau_n) + (s+2-t-1/t)f_{m-1}g_{m-1}\tau_n.$$

Proof. By [17] (see also [8, p.309]), the Riley polynomial is $\phi_K(s,t) = z_{1,1} + (1-t)z_{1,2}$, where $z_{i,j}$ is the (i,j)-entry of W^n. From Lemma 3.5 and the recursive formula for τ_k,

$$\begin{aligned} \phi_K(s,t) &= (w_{1,1}\tau_n - \tau_{n-1}) + (1-t)w_{1,2}\tau_n \\ &= \tau_{n+1} - (\mathrm{tr}W)\tau_n + (w_{1,1} + (1-t)w_{1,2})\tau_n \\ &= (\tau_{n+1} - \tau_n) + (w_{1,1} + (1-t)w_{1,2} + 1 - \mathrm{tr}W)\tau_n. \end{aligned}$$

By Proposition 3.1, $\mathrm{tr}W = f_m^2 + f_{m-1}^2 - s(t+1/t)g_{m-1}^2$. Thus we have

$$w_{1,1} + (1-t)w_{1,2} + 1 - \mathrm{tr}W = (2-t-1/t)f_{m-1}g_{m-1} + 1 - f_{m-1}^2 + sg_{m-1}^2.$$

Since

$$\begin{aligned} 1 - f_{m-1}^2 + sg_{m-1}^2 &= -sg_{m-1}g_{m-2} + sg_{m-1}^2 \\ &= sg_{m-1}(g_{m-1} - g_{m-2}) \\ &= sg_{m-1}f_{m-1}, \end{aligned}$$

we obtain $\phi_K(s,t) = (\tau_{n+1} - \tau_n) + (s+2-t-1/t)f_{m-1}g_{m-1}\tau_n$. □

For convenience, we introduce a variable $T = t+1/t$. Then the Riley polynomial of K is expressed as $\phi_K(s,T) = (\tau_{n+1} - \tau_n) + (s+2-T)f_{m-1}g_{m-1}\tau_n$.

We remark that the Riley polynomial of K is also described by [15] in a different form.

Example 3.7. If $n = 1$, then

$$\begin{aligned} \phi_K(s,T) &= (\tau_2 - \tau_1) + (s+2-T)f_{m-1}g_{m-1}\tau_1 \\ &= (\mathrm{tr}W - 1) + (s+2-T)f_{m-1}g_{m-1} \\ &= s(s+2-T)g_{m-1}^2 + 1 + (s+2-T)f_{m-1}g_{m-1} \\ &= (s+2-T)g_{m-1}(sg_{m-1} + f_{m-1}) + 1 \\ &= (s+2-T)g_{m-1}f_m + 1. \end{aligned}$$

Thus Riley's equation $\phi_K(s,T) = 0$ has the unique solution $T = s+2+1/(f_m g_{m-1})$ for any $s > 0$. If $m > 0$, then $T > s+2 > 2$, because $f_m > 0$ and $g_{m-1} > 0$. Hence we have a real solution $t = (T + \sqrt{T^2-4})/2 > 1$. By Lemma 2.4(1), we also have $s+2 < T < s+2+4/(sg_{m-1}^2)$.

If $n = -1$, then $\phi_K(s,T) = 1 + (T-s-2)f_{m-1}g_{m-1}$. Hence the equation $\phi_K(s,T) = 0$ also has the unique solution $T = s+2-1/(f_{m-1}g_{m-1})$. If $m < 0$, then $T > s+2 > 2$, so there exists a real solution $t > 1$ as above. Again, $s+2 < T < s+2+4/(sg_{m-1}^2)$ by Lemma 2.4(2).

4. Solutions of Riley's equation

In this section, we examine when Riley's equation $\phi_K(s,T) = 0$ has a pair of real solutions (s,T). In fact, we can choose T satisfying $s+2+c/s < T < s+2+d/s$, where c and d are constants depending only n, for any $s > 0$, unless $n = \pm 1$.

Let k be a positive integer. For $z = e^{i\theta}$ $(0 \le \theta \le \pi)$, set $\mathcal{T}_k(z) = z^{k-1} + z^{k-3} + \cdots + z^{3-k} + z^{1-k}$. If $z \ne \pm 1$, then $\mathcal{T}_k(z) = (z^k - z^{-k})/(z - z^{-1})$. Define $\mathcal{T}_0 = 0$ and $\mathcal{T}_{-k}(z) = -\mathcal{T}_k(z)$. Since $\mathcal{T}_k(z)$ is symmetric for z and z^{-1}, it can be expanded as a polynomial of $z + z^{-1}$. Furthermore, a recursive relation

$$\mathcal{T}_{k+1}(z) - (z + z^{-1})\mathcal{T}_k(z) + \mathcal{T}_{k-1}(z) = 0$$

holds. Also, $\mathcal{T}_k(1) = k$ and $\mathcal{T}_k(-1) = (-1)^{k-1}k$ for any integer k.

Lemma 4.1. (1) *Let* $k \geq 1$. *Then,* $\mathcal{T}_k(e^{\frac{\pi}{2k+1}i}) = \mathcal{T}_{k+1}(e^{\frac{\pi}{2k+1}i})$. *This value is positive.*

(2) *Let* $m \geq 2$. *Then,* $\mathcal{T}_k(e^{\frac{3\pi}{2k+1}i}) = \mathcal{T}_{k+1}(e^{\frac{3\pi}{2k+1}i})$. *This value is negative.*

Proof. (1) Let $z = e^{\frac{\pi}{2k+1}i}$. Then the fact that $z^{2k+1} = -1$ immediately implies $\mathcal{T}_k(z) = \mathcal{T}_{k+1}(z)$. A direct calculation shows

$$\mathcal{T}_k(z) = \frac{\sin\frac{k\pi}{2k+1}}{\sin\frac{\pi}{2k+1}} > 0.$$

(2) Similarly, set $z = e^{\frac{3\pi}{2k+1}i}$. Then $z^{2k+1} = -1$ holds again. Hence we have $\mathcal{T}_k(z) = \mathcal{T}_{k+1}(z)$, and

$$\mathcal{T}_k(z) = \frac{\sin\frac{3k\pi}{2k+1}}{\sin\frac{3\pi}{2k+1}} < 0.$$

□

Now, fix an $s > 0$. We introduce a function $\Phi : [s+2, s+2+4/(sg_{m-1}^2)] \to \mathbb{R}$ by

$$\Phi(T) = (\mathcal{T}_{n+1}(z) - \mathcal{T}_n(z)) + (s+2-T)f_{m-1}g_{m-1}\mathcal{T}_n(z), \tag{4.1}$$

where $z = (\tau + i\sqrt{4-\tau^2})/2$ with $\tau = s(2+s-T)g_{m-1}^2 + 2$.

Note that $-2 \leq \tau \leq 2$. We will seek a solution T for $\Phi(T) = 0$ satisfying $s+2 < T < s+2+4/(sg_{m-1}^2)$, because it gives a pair of solutions (s,T) for Riley's equation $\phi_K(s,T) = 0$.

Proposition 4.2. *Suppose* $n \neq \pm 1$. *For any* $s > 0$, *Riley's equation* $\phi_K(s,T) = 0$ *has a solution* T *satisfying* $s+2+c/(sg_{m-1}^2) < T < s+2+d/(sg_{m-1}^2)$, *where* c *and* d *are constants in* $(0,4)$ *depending only on* n. *In particular,* $\phi_K(s,t) = 0$ *has a solution* $t > 1$ *for any* $s > 0$.

Proof. Suppose $n > 1$. By Lemma 4.1,

$$\mathcal{T}_{n+1}(e^{\frac{\pi}{2n+1}i}) = \mathcal{T}_n(e^{\frac{\pi}{2n+1}i}), \quad \mathcal{T}_{n+1}(e^{\frac{3\pi}{2n+1}i}) = \mathcal{T}_n(e^{\frac{3\pi}{2n+1}i}).$$

Let $c = 2 - 2\cos\frac{\pi}{2n+1}$ and $c' = 2 - 2\cos\frac{3\pi}{2n+1}$. Then $c, c' \in (0,4)$ and $c < c'$.

If $T = s+2+c/(sg_{m-1}^2)$ (resp. $s+2+c'/(sg_{m-1}^2)$ then $\tau = 2-c$ (resp. $2-c'$). Thus

$$\begin{aligned}
\Phi(s+2+\frac{c}{sg_{m-1}^2}) &= -\frac{cf_{m-1}}{sg_{m-1}}\mathcal{T}_n(e^{\frac{\pi}{2n+1}i}),\\
\Phi(s+2+\frac{c'}{sg_{m-1}^2}) &= -\frac{c'f_{m-1}}{sg_{m-1}}\mathcal{T}_n(e^{\frac{3\pi}{2n+1}i}).
\end{aligned}$$

By Lemma 4.1, these values have distinct signs. We remark that $\Phi(T)$ is a polynomial function of T, so it is continuous. Thus if $n > 1$, we have a solution T for $\Phi(T) = 0$, satisfying $s+2+c/(sg_{m-1}^2) < T < s+2+c'/(sg_{m-1}^2)$, from the

Intermediate-Value Theorem. Since $T > 2$, $t + 1/t = T$ has a real solution for t. If we choose $t = (T + \sqrt{T^2 - 4})/2$, then $t > 1$.

Suppose $n < -1$. Set $l = |n|$. If $l > 2$, then set $d = 2 - 2\cos\frac{\pi}{2l-1}$ and $d' = 2 - 2\cos\frac{3\pi}{2l-1}$. Then $d, d' \in (0, 4)$ and $d < d'$. As before,

$$\mathcal{T}_{l-1}(e^{\frac{\pi}{2l-1}i}) = \mathcal{T}_l(e^{\frac{\pi}{2l-1}i}), \quad \mathcal{T}_{l-1}(e^{\frac{3\pi}{2l-1}i}) = \mathcal{T}_l(e^{\frac{3\pi}{2l-1}i}).$$

by Lemma 4.1. Thus

$$\begin{aligned} \Phi(s + 2 + \frac{d}{sg_{m-1}^2}) &= \frac{d f_{m-1}}{sg_{m-1}^2}\mathcal{T}_l(e^{\frac{\pi}{2l-1}i}), \\ \Phi(s + 2 + \frac{d'}{sg_{m-1}^2}) &= \frac{d' f_{m-1}}{sg_{m-1}^2}\mathcal{T}_l(e^{\frac{3\pi}{2l-1}i}). \end{aligned}$$

Since these values have distinct signs, we have a solution T with $s+2+d/(sg_{m-1}^2) < T < s + 2 + d'/(sg_{m-1}^2)$, if $l > 2$, as before.

When $l = 2$, we have

$$\Phi(s + 2 + \frac{1}{sg_{m-1}^2}) = \frac{f_{m-1}}{sg_{m-1}}, \quad \Phi(s + 2 + \frac{2}{sg_{m-1}^2}) = -1,$$

$$\Phi(s + 2 + \frac{3}{sg_{m-1}^2}) = -2 - \frac{3f_{m-1}}{sg_{m-1}}.$$

If $m > 0$, then $\Phi(s + 2 + 1/(sg_{m-1}^2)) > 0$, since $f_{m-1}, g_{m-1} > 0$. Therefore there exists a solution T with $s + 2 + 1/(sg_{m-1}^2) < T < s + 2 + 2/(sg_{m-1}^2)$. If $m < 0$, $\Phi(s + 2 + 3/(sg_{m-1}^2)) > 0$, since $3f_{m-1} > -2sg_{m-1}$ by Lemma 2.4(3). Thus there exists a solution T with $s + 2 + 2/(sg_{m-1}^2) < T < s + 2 + 3/(sg_{m-1}^2)$ □

5. Longitudes

For $s > 0$, let $\rho_s : G \to SL_2(\mathbb{R})$ be the representation defined by the correspondence

$$\rho_s(x) = \begin{pmatrix} \sqrt{t} & 0 \\ 0 & \frac{1}{\sqrt{t}} \end{pmatrix}, \quad \rho_s(y) = \begin{pmatrix} \frac{t-s-1}{\sqrt{t}-\frac{1}{\sqrt{t}}} & \frac{s}{(\sqrt{t}-\frac{1}{\sqrt{t}})^2} - 1 \\ -s & \frac{s+1-\frac{1}{t}}{\sqrt{t}-\frac{1}{\sqrt{t}}} \end{pmatrix}. \tag{5.1}$$

For $Q = \begin{pmatrix} t-1 & 1 \\ 0 & \sqrt{t} - \frac{1}{\sqrt{t}} \end{pmatrix}$,

$$Q^{-1}\rho_s(x)Q = \begin{pmatrix} \sqrt{t} & \frac{1}{\sqrt{t}} \\ 0 & \frac{1}{\sqrt{t}} \end{pmatrix}, \quad Q^{-1}\rho_s(y)Q = \begin{pmatrix} \sqrt{t} & 0 \\ -s\sqrt{t} & \frac{1}{\sqrt{t}} \end{pmatrix}.$$

Therefore, ρ_s is conjugate with ρ defined in Section 3. This implies that if s and t satisfy Riley's equation $\phi_K(s, t) = 0$ then ρ_s gives a representation of G as well as ρ. In this section, we examine the image of the longitude $\mathcal{L}$ of G under ρ_s.

Throughout the section, let $U = \rho_s(w)$ and $u_{i,j}$ be its (i, j)-entry, and let $v_{i,j}$ be the entries of U^n. Also, set $\sigma = \frac{s(\sqrt{t}-\frac{1}{\sqrt{t}})^2}{(\sqrt{t}-\frac{1}{\sqrt{t}})^2-s}$.

Lemma 5.1. *For* w_*^n*, we have* $\rho_s(w_*^n) = \begin{pmatrix} v_{1,1} & \frac{v_{2,1}}{\sigma} \\ v_{1,2}\sigma & v_{2,2} \end{pmatrix}$.

Proof. By a direct calculation,

$$\rho_s(xy^{-1}) = \begin{pmatrix} \frac{t-1+st}{t-1} & \frac{s\sqrt{t}}{\sigma} \\ \frac{s}{\sqrt{t}} & \frac{t-1-s}{t-1} \end{pmatrix}, \quad \rho_s(y^{-1}x) = \begin{pmatrix} \frac{t-1+st}{t-1} & \frac{s}{\sqrt{t}\sigma} \\ s\sqrt{t} & \frac{t-1-s}{t-1} \end{pmatrix},$$

$$\rho_s(x^{-1}y) = \begin{pmatrix} \frac{t-1-s}{t-1} & -\frac{s}{\sqrt{t}\sigma} \\ -s\sqrt{t} & \frac{t-1+st}{t-1} \end{pmatrix}, \quad \rho_s(yx^{-1}) = \begin{pmatrix} \frac{t-1-s}{t-1} & -\frac{s\sqrt{t}}{\sigma} \\ -\frac{s}{\sqrt{t}} & \frac{t-1+st}{t-1} \end{pmatrix}.$$

Thus we see that the $(1,2)$-entry of $\rho_s(y^{-1}x)$ is the $(2,1)$-entry of $\rho_s(xy^{-1})$ divided by σ, the $(2,1)$-entry of $\rho_s(y^{-1}x)$ is the $(1,2)$-entry of $\rho_s(xy^{-1})$ multiplied by σ, and the others of $\rho_s(y^{-1}x)$ coincide with those of $\rho_s(xy^{-1})$. The same relation between entries holds for $\rho_s(x^{-1}y)$ and $\rho_s(yx^{-1})$.

In general, such a relation is preserved under the matrix multiplication;

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}\begin{pmatrix} p & q \\ r & s \end{pmatrix} = \begin{pmatrix} ap+br & aq+bs \\ cp+dr & cq+ds \end{pmatrix},$$

$$\begin{pmatrix} p & \frac{r}{\sigma} \\ q\sigma & s \end{pmatrix}\begin{pmatrix} a & \frac{c}{\sigma} \\ b\sigma & d \end{pmatrix} = \begin{pmatrix} ap+br & \frac{cp+dr}{\sigma} \\ (aq+bs)\sigma & cq+ds \end{pmatrix}.$$

Thus we can confirm that the same relation holds for $\rho_s(w^n)$ and $\rho_s(w_*^n)$. □

Proposition 5.2. *For the longitude* $\mathcal{L}$ *of* G*, the matrix* $\rho_s(\mathcal{L})$ *is diagonal, and the* $(1,1)$*-entry of* $\rho_s(\mathcal{L})$ *is a positive real number.*

Proof. The first assertion follows from the facts that for a meridian x, $\rho_s(x)$ is diagonal but $\rho_s(x) \neq \pm I$ and that x and $\mathcal{L}$ commute.

Since $\mathcal{L} = w_*^n w^n$, Lemma 5.1 implies that

$$\begin{aligned} \rho_s(\mathcal{L}) = \rho_s(w_*^n)\rho_s(w^n) &= \begin{pmatrix} v_{1,1} & \frac{v_{2,1}}{\sigma} \\ v_{1,2}\sigma & v_{2,2} \end{pmatrix}\begin{pmatrix} v_{1,1} & v_{1,2} \\ v_{2,1} & v_{2,2} \end{pmatrix} \\ &= \begin{pmatrix} v_{1,1}^2 + \frac{v_{2,1}^2}{\sigma} & v_{1,1}v_{1,2} + \frac{v_{2,1}v_{2,2}}{\sigma} \\ v_{1,1}v_{1,2}\sigma + v_{2,1}v_{2,2} & v_{1,2}^2\sigma + v_{2,2}^2 \end{pmatrix}. \end{aligned}$$

Hence the $(1,1)$-entry is $v_{1,1}^2 + v_{2,1}^2/\sigma$, which is positive, because $s > 0$ and $(\sqrt{t} - 1/\sqrt{t})^2 - s = T - s - 2 > 0$ from Proposition 4.2. □

Remark 5.3. Since $\rho_s(\mathcal{L})$ is diagonal, we also obtain an equation $v_{1,1}v_{1,2}\sigma + v_{2,1}v_{2,2} = 0$. This will be used in the proof of Lemma 5.6.

For $W = \rho(w)$, recall that $w_{i,j}$ is its entry.

Lemma 5.4. *For $U = \rho_s(w)$,*

$$\begin{aligned} u_{1,1} &= w_{1,1} + \frac{w_{2,1}}{t-1}, \quad u_{1,2} = \sqrt{t}\left(w_{1,2} - \frac{w_{1,1}}{t-1}\right) + \frac{\sqrt{t}}{t-1}\left(w_{2,2} - \frac{w_{2,1}}{t-1}\right), \\ u_{2,1} &= \frac{w_{2,1}}{\sqrt{t}}, \quad u_{2,2} = w_{2,2} - \frac{w_{2,1}}{t-1}. \end{aligned}$$

Proof. This immediately follows by calculating the product $U = QWQ^{-1}$. □

By Lemma 3.5, we have

$$W^n = \begin{pmatrix} w_{1,1}\tau_n - \tau_{n-1} & w_{1,2}\tau_n \\ w_{2,1}\tau_n & \tau_{n+1} - w_{1,1}\tau_n \end{pmatrix}.$$

Lemma 5.5. *For $U^n = \rho_s(w^n)$, we have*

$$\begin{aligned} v_{1,1} &= u_{1,1}\tau_n - \tau_{n-1}, \quad v_{1,2} = u_{1,2}\tau_n, \\ v_{2,1} &= u_{2,1}\tau_n \quad v_{2,2} = \tau_{n+1} - u_{1,1}\tau_n. \end{aligned}$$

Proof. Calculate the product $U^n = QW^nQ^{-1}$. Then $v_{1,1} = (w_{1,1}+w_{2,1}/(t-1))\tau_n - \tau_{n-1} = u_{1,1}\tau_n - \tau_{n-1}$ by Lemma 5.4. For $v_{1,2}$, we have

$$\begin{aligned} v_{1,2} = \frac{\sqrt{t}}{(t-1)^2}\Big(&(t-1)\tau_{n-1} - (t-1)w_{1,1}\tau_n + (t-1)^2 w_{1,2}\tau_n \\ &+ (t-1)\tau_{n+1} - (t-1)w_{1,1}\tau_n - w_{2,1}\tau_n\Big). \end{aligned}$$

Recall that the Riley polynomial is $\phi_K(s,t) = (w_{1,1}\tau_n - \tau_{n-1}) + (1-t)w_{1,2}\tau_n$. (See the proof of Proposition 3.6.) Since $\phi_K(s,t) = 0$, $\tau_{n-1} = w_{1,1}\tau_n + (1-t)w_{1,2}\tau_n$. Hence

$$\begin{aligned} v_{1,2} &= \frac{\sqrt{t}}{(t-1)^2}\Big((t-1)\tau_{n+1} - (w_{2,1} + (t-1)w_{1,1})\tau_n\Big) \\ &= \frac{\sqrt{t}}{(t-1)^2}\Big(\big((t-1)w_{2,2} - w_{2,1}\big)\tau_n - (t-1)\tau_{n-1}\Big) \\ &= \frac{\sqrt{t}}{t-1}\Big((w_{2,2} - \frac{w_{2,1}}{t-1})\tau_n - \tau_{n-1}\Big) \\ &= \frac{\sqrt{t}}{t-1}(u_{2,2}\tau_n - \tau_{n-1}), \end{aligned}$$

by using the recursion $\tau_{n+1} - (w_{1,1} + w_{2,2})\tau_n + \tau_{n-1} = 0$. By Lemma 5.4,

$$u_{2,2} = \frac{t-1}{\sqrt{t}}\left(u_{1,2} - \sqrt{t}\left(w_{1,2} - \frac{w_{1,1}}{t-1}\right)\right).$$

Substituting this and $\phi_K(s,t) = 0$,

$$\begin{aligned} v_{1,2} &= u_{1,2}\tau_n + \frac{\sqrt{t}}{t-1}\Big(w_{1,1}\tau_n - \tau_{n-1} + (1-t)w_{1,2}\tau_n\Big) \\ &= u_{1,2}\tau_n. \end{aligned}$$

It is straightforward to check $v_{2,1}$ and $v_{2,2}$. We omit them. □

Let B_s be the $(1,1)$-entry of the matrix $\rho_s(\mathcal{L})$.

Lemma 5.6. *$B_s = -u_{2,1}/(u_{1,2}\sigma)$.*

Proof. As noted in Remark 5.3, $v_{1,1}v_{1,2}\sigma + v_{2,1}v_{2,2} = 0$. Since $\det U^n = v_{1,1}v_{2,2} - v_{1,2}v_{2,1} = 1$, we have

$$\begin{aligned} v_{1,2}B_s &= v_{1,1}^2 v_{1,2} + \frac{v_{1,2}v_{2,1}^2}{\sigma} \\ &= v_{1,1}^2 v_{1,2} + \frac{v_{2,1}}{\sigma}(v_{1,1}v_{2,2} - 1) \\ &= v_{1,1}^2 v_{1,2} + \frac{v_{1,1}}{\sigma}(-v_{1,1}v_{1,2}\sigma) - \frac{v_{2,1}}{\sigma} \\ &= -\frac{v_{2,1}}{\sigma}. \end{aligned}$$

Assume $v_{1,2} = 0$. Then $v_{2,1} = 0$. By Lemmas 5.4 and 5.5, $v_{2,1} = u_{2,1}\tau_n = w_{2,1}\tau_n/\sqrt{t}$. Since $w_{2,1} \neq 0$ by Lemma 3.4, we have $\tau_n = 0$. Recall that $\phi_K(s,t) = (\tau_{n+1} - \tau_n) + (s + 2 - t - 1/t)f_{m-1}g_{m-1}\tau_n$ is zero. Thus $\tau_{n+1} = 0$. Then the recursive formula for τ_k implies $\tau_{n-1} = 0$. In turn, all $\tau_k = 0$. But this is impossible, because $\tau_1 = 1$. Hence $v_{1,2} \neq 0$, so $B_s = -v_{2,1}/(v_{1,2}\sigma)$.

By Lemma 5.5, $v_{1,2} = u_{1,2}\tau_n$ and $v_{2,1} = u_{2,1}\tau_n$. Thus we have shown that $B_s = -u_{2,1}/(u_{1,2}\sigma)$. □

Proposition 5.7. *For the longitude $\mathcal{L}$, the $(1,1)$-entry B_s of $\rho_s(\mathcal{L})$ is given as*

$$B_s = \frac{-f_m + tf_{m-1}}{-f_{m-1} + tf_m}. \tag{5.2}$$

Proof. From Lemma 5.6, $B_s = -u_{2,1}/(u_{1,2}\sigma)$. Then Lemma 5.4 and Proposition 3.1,

$$\begin{aligned} u_{1,2} &= \frac{\sqrt{t}}{t-1}\left(f_{m-1}^2 - f_m^2 + (st - \frac{s}{t})g_{m-1}^2\right) - \frac{(t-1)^2 + st}{\sqrt{t}(t-1)^2}g_{m-1}(f_m - tf_{m-1}), \\ u_{2,1} &= \frac{s}{\sqrt{t}}g_{m-1}(f_m - tf_{m-1}). \end{aligned}$$

For the first term of $u_{1,2}$, Lemma 2.3 implies

$$\begin{aligned} f_{m-1}^2 - f_m^2 + (st - \frac{s}{t})g_{m-1}^2 &= (f_{m-1} - f_m)(f_{m-1} + f_m) + (t - \frac{1}{t})sg_{m-1}^2 \\ &= -sg_{m-1}(f_{m-1} + f_m) + (t - \frac{1}{t})sg_{m-1}^2 \\ &= sg_{m-1}\left(-f_{m-1} - f_m + (t - \frac{1}{t})g_{m-1}\right) \\ &= sg_{m-1}\left(-f_{m-1} - f_m + (t - \frac{1}{t})\frac{f_m - f_{m-1}}{s}\right) \\ &= sg_{m-1}\left(\frac{t^2 - 1 - st}{st}f_m - \frac{t^2 - 1 + st}{st}f_{m-1}\right). \end{aligned}$$

Thus, dividing $u_{1,2}$ by $sg_{m-1}/\sqrt{t}$ gives

$$\frac{t}{t-1}\left(\frac{t^2-1-st}{st}f_m-\frac{t^2-1+st}{st}f_{m-1}\right)-\frac{(t-1)^2+st}{s(t-1)^2}(f_m-tf_{m-1}).$$

The coefficient of f_m is t/σ, and that of f_{m-1} is $-1/\sigma$. Hence we have

$$B_s=-\frac{u_{2,1}}{u_{1,2}\sigma}=\frac{-f_m+tf_{m-1}}{-f_{m-1}+tf_m}.$$

□

6. Limits

Let $r=p/q$ be a rational number, and let $K(r)$ denote the resulting manifold by r-surgery on K. In other words, $K(r)$ is obtained by attaching a solid torus V to the knot exterior $E(K)$ along their boundaries so that the loop $x^p\mathcal{L}^q$ bounds a meridian disk of V, where x and $\mathcal{L}$ are a meridian and longitude of K.

Our representation $\rho_s : G \to SL_2(\mathbb{R})$ induces a homomorphism $\pi_1(K(r)) \to SL_2(\mathbb{R})$ if and only if $\rho_s(x)^p\rho_s(\mathcal{L})^q=I$. Since both of $\rho_s(x)$ and $\rho_s(\mathcal{L})$ are diagonal (see (5.1) and Proposition 5.2), this is equivalent to the single equation

$$A_s^pB_s^q=1, \tag{6.1}$$

where A_s and B_s are the $(1,1)$-entries of $\rho_s(x)$ and $\rho_s(\mathcal{L})$, respectively. We remark that $A_s=\sqrt{t}$ (>1) is a positive real number, so is B_s by Proposition 5.2. Hence the equation (6.1) is furthermore equivalent to the equation

$$-\frac{\log B_s}{\log A_s}=\frac{p}{q}. \tag{6.2}$$

Let $g:(0,\infty)\to\mathbb{R}$ be a function defined by

$$g(s)=-\frac{\log B_s}{\log A_s}.$$

We will examine the image of g.

Lemma 6.1. (1)

$$\lim_{s\to+0}t=\begin{cases}\infty & \textit{if } n\neq\pm1,\\ \frac{2m+1+\sqrt{4m+1}}{2m} & \textit{if } n=1,\\ \frac{2m-1+\sqrt{1-4m}}{2m} & \textit{if } n=-1.\end{cases}$$

(2) $\lim_{s\to\infty}t=\infty$.
(3) $\lim_{s\to\infty}(t-s)=2$.
(4) $\lim_{s\to\infty}\dfrac{t}{s}=1$.

Proof. (1) If $n = 1$, then $T = s + 2 + 1/(f_m g_{m-1})$ is the unique solution for $\phi_K(s, T) = 0$ (see Example 3.7). From Lemma 2.2, we have $\lim_{s\to+0} f_m = 1$ and $\lim_{s\to+0} g_{m-1} = m$. Hence $\lim_{s\to+0} T = 2 + 1/m$. (Recall $m > 0$ when $n = 1$.) Since $t = (T + \sqrt{T^2 - 4})/2$, $\lim_{s\to+0} t = (2m + 1 + \sqrt{4m + 1})/(2m)$.

If $n = -1$, then $T = s + 2 - 1/(f_{m-1} g_{m-1})$ by Example 3.7. Recall $m < 0$ when $n = -1$. Then $\lim_{s\to+0} f_{m-1} = 1$ and $\lim_{s\to+0} g_{m-1} = m$. Thus $\lim_{s\to+0} T = 2 - 1/m$, so $\lim_{s\to+0} t = (2m - 1 + \sqrt{1 - 4m})/(2m)$.

Assume $n \neq \pm 1$. From Proposition 4.2, we have $s + 2 + c/(s g_{m-1}^2) < T$, where c is a positive constant. Hence $\lim_{s\to+0} T = \lim_{s\to+0} t = \infty$.

(2) As $T > s + 2$, $\lim_{s\to\infty} T = \lim_{s\to\infty} t = \infty$.

(3) Since $s + 2 < t + 1/t < s + 2 + 4/(s g_{m-1}^2)$, (2) implies $\lim_{s\to\infty}(t - s) = 2$.

(4) From $s + 2 < T < s + 2 + 4/(s g_{m-1}^2)$ again, we have $\lim_{s\to\infty} T/s = 1$, which implies $\lim_{s\to\infty} t/s = 1$ □

Let $F_k = t^{k-1}(-f_k + t f_{k-1})$ for any integer k.

Lemma 6.2. *If* $0 < k \leq |m|$, *then* $\lim_{s\to\infty} F_k = 1$.

Proof. First, $F_1 = -f_1 + t f_0 = t - s - 1$. Hence $\lim_{s\to\infty} F_1 = 1$ by Lemma 6.1(3).

Let $n \neq \pm 1$. By Proposition 4.2, we have

$$s + 2 + \frac{c}{s g_{m-1}^2} < t + \frac{1}{t} < s + 2 + \frac{d}{s g_{m-1}^2}, \tag{6.3}$$

where c and d are constants depending on only n. Multiplying $t^{k-1} f_{k-1}$ to (6.3) gives

$$(s+2) f_{k-1} t^{k-1} + \frac{c t^{k-1} f_{k-1}}{s g_{m-1}^2} < t^k f_{k-1} + t^{k-2} f_{k-1} < (s+2) f_{k-1} t^{k-1} + \frac{d t^{k-1} f_{k-1}}{s g_{m-1}^2}.$$

Since $(s + 2) f_{k-1} = f_k + f_{k-2}$ by the recursion,

$$\begin{aligned} (t^{k-1} f_{k-2} - t^{k-2} f_{k-1}) + \frac{c t^{k-1} f_{k-1}}{s g_{m-1}^2} &< t^k f_{k-1} - t^{k-1} f_k \\ &< (t^{k-1} f_{k-2} - t^{k-2} f_{k-1}) + \frac{d t^{k-1} f_{k-1}}{s g_{m-1}^2}. \end{aligned}$$

Then we have

$$F_{k-1} + \frac{c t^{k-1} f_{k-1}}{s g_{m-1}^2} < F_k < F_{k-1} + \frac{d t^{k-1} f_{k-1}}{s g_{m-1}^2}. \tag{6.4}$$

The degree of f_{k-1} is $k - 1$, but that of g_{m-1} is $|m| - 1$. Hence Lemma 6.1(4) implies

$$\lim_{s\to\infty} \frac{c t^{k-1} f_{k-1}}{s g_{m-1}^2} = \lim_{s\to\infty} \frac{d t^{k-1} f_{k-1}}{s g_{m-1}^2} = 0$$

as long as $k \leq |m|$. Thus $\lim_{s\to\infty} F_k = 1$ will follow from $\lim_{s\to\infty} F_{k-1} = 1$.

Suppose $n = 1$. Riley's equation has the unique solution

$$t + \frac{1}{t} = s + 2 + \frac{1}{f_m g_{m-1}} \tag{6.5}$$

(see Example 3.7).

Multiplying $t^{k-1}f_{k-1}$ to (6.5) gives

$$t^k f_{k-1} + t^{k-2} f_{k-1} = (s+2)f_{k-1}t^{k-1} + \frac{t^{k-1}f_{k-1}}{f_m g_{m-1}}.$$

Since $(s+2)f_{k-1} = f_k + f_{k-2}$,

$$t^k f_{k-1} - t^{k-1} f_k = t^{k-1} f_{k-2} - t^{k-2} f_{k-1} + \frac{t^{k-1}f_{k-1}}{f_m g_{m-1}}.$$

That is, $F_k = F_{k-1} + t^{k-1}f_{k-1}/(f_m g_{m-1})$. Hence the fact that $\lim_{s\to\infty} F_{k-1} = 1$ and $\lim_{s\to\infty} t^{k-1}f_{k-1}/(f_m g_{m-1}) = 0$ imply $\lim_{s\to\infty} F_k = 1$.

Finally, suppose $n = -1$. Riley's equation has the unique solution

$$t + \frac{1}{t} = s + 2 - \frac{1}{f_{m-1}g_{m-1}}$$

as in Example 3.7. Multiplying $t^{k-1}f_{k-1}$ gives $F_k = F_{k-1} - t^{k-1}f_{k-1}/(f_{m-1}g_{m-1})$. Since $m < 0$, f_{m-1} and g_{m-1} have degree $|m|$ and $|m|-1$, respectively. Thus the fact that $\lim_{s\to\infty} F_{k-1} = 1$ and $\lim_{s\to\infty} t^{k-1}f_{k-1}/(f_{m-1}g_{m-1}) = 0$ imply $\lim_{s\to\infty} F_k = 1$, again. □

Lemma 6.3. (1) $\lim\limits_{s\to+0} B_s = 1$.

(2) $\lim\limits_{s\to\infty} B_s t^{2m} = 1$.

Proof. (1) By Proposition 5.7,

$$B_s = \frac{-f_m + tf_{m-1}}{-f_{m-1} + tf_m}.$$

By Lemma 2.2, $\lim_{s\to+0} f_m = \lim_{s\to+0} f_{m-1} = 1$. Thus Lemma 6.1(1) implies $\lim_{s\to+0} B_s = 1$.

(2) Let $m > 0$. We decompose $B_s t^{2m}$ as

$$\begin{aligned} B_s t^{2m} &= t^{m-1}(-f_m + tf_{m-1}) \cdot \frac{t^{m+1}}{-f_{m-1} + tf_m} \\ &= F_m \cdot \frac{t^{m+1}}{-f_{m-1} + tf_m}. \end{aligned}$$

Since the degree of f_k $(k > 0)$ is k and f_m is monic,

$$\lim_{s\to\infty} \frac{t^{m+1}}{-f_{m-1} + tf_m} = 1.$$

Then we have $\lim_{s\to\infty} B_s t^{2m} = 1$ by combined with Lemma 6.2.

Let $m < 0$. Set $l = -m > 0$. Recall $f_m = f_{l-1}$ and $f_{m-1} = f_l$. We decompose $B_s t^{2m}$ as

$$B_s t^{2m} = \frac{-f_{l-1} + tf_l}{t^{l+1}} \cdot \frac{1}{t^{l-1}(-f_l + tf_{l-1})} = \frac{-f_{l-1} + tf_l}{t^{l+1}} \cdot \frac{1}{F_l}.$$

As before,

$$\lim_{s\to\infty} \frac{-f_{l-1} + tf_l}{t^{l+1}} = 1 \text{ and } \lim_{s\to\infty} F_l = 1.$$

Thus $\lim_{s\to\infty} B_s t^{2m} = 1$ again. □

Proposition 6.4. *The image of g contains an open interval $(0, 4m)$ (resp. $(4m, 0)$) if $m > 0$ (resp. $m < 0$).*

Proof. By Lemma 6.3(1), $\lim_{s\to+0} \log B_s = 0$. Hence

$$\lim_{s\to+0} g(s) = -\lim_{s\to+0} \frac{\log B_s}{\log A_s} = -\lim_{s\to+0} \frac{\log B_s}{\log\sqrt{t}} = 0.$$

Also, we have $\lim_{s\to\infty}(\log B_s + 2m\log t) = 0$ by Lemma 6.3(2). Thus

$$\lim_{s\to\infty} g(s) = -\lim_{s\to\infty} \frac{\log B_s}{\log A_s} = -\lim_{s\to\infty} \frac{2\log B_s}{\log t} = 4m.$$

Hence the image of g contains an interval $(0, 4m)$ or $(4m, 0)$, according as the sign of m. □

7. Proof of Theorem

The universal covering group $\widetilde{SL_2(\mathbb{R})}$ of $SL_2(\mathbb{R})$ can be described as

$$\widetilde{SL_2(\mathbb{R})} = \{(\gamma, \omega) \mid \gamma \in \mathbb{C}, |\gamma| < 1, -\infty < \omega < \infty\},$$

(see [1]). Let $\chi : \widetilde{SL_2(\mathbb{R})} \to SL_2(\mathbb{R})$ be the covering projection. Then $\ker\chi = \{(0, 2m\pi) \mid m \in \mathbb{Z}\}$ is isomorphic to $\mathbb{Z}$.

Since the knot exterior $E(K)$ of K satisfies $H^2(E(K); \mathbb{Z}) = 0$, any $\rho_s : G \to SL_2(\mathbb{R})$ lifts to a representation $\tilde{\rho} : G \to \widetilde{SL_2(\mathbb{R})}$ ([9]). Moreover, any two lifts $\tilde{\rho}$ and $\tilde{\rho}'$ are related as follows:

$$\tilde{\rho}'(g) = h(g)\tilde{\rho}(g),$$

where $h : G \to \ker\chi \subset \widetilde{SL_2(\mathbb{R})}$. Since $\ker\chi$ is abelian, the homomorphism h factors through $H_1(E(K))$, so it is determined only by the value $h(x)$ of a meridian x (see [14]).

Lemma 7.1. *Let $\tilde{\rho} : G \to \widetilde{SL_2(\mathbb{R})}$ be a lift of ρ_s. Then replacing $\tilde{\rho}$ by a representation $\tilde{\rho}' = h \cdot \tilde{\rho}$ for some $h : G \to \widetilde{SL_2(\mathbb{R})}$, we can suppose that $\tilde{\rho}(\pi_1(\partial E(K)))$ is contained in the subgroup $(-1, 1) \times \{0\}$ of $\widetilde{SL_2(\mathbb{R})}$.*

Proof. This is proved in [11, Section 7] for twist knots. Since our knot K has genus one, the argument works without any change. □

Proof of Theorem 1.1. Suppose that $m, n > 0$. Let $r = p/q \in (0, 4m)$. By Proposition 6.4, we can find s so that $g(s) = r$. Choose a lift $\tilde{\rho}$ of ρ_s so that $\tilde{\rho}(\pi_1(\partial E(K))) \subset (-1, 1) \times \{0\}$ (Lemma 7.1). Then $\rho_s(x^p\mathcal{L}^q) = I$, so $\chi(\tilde{\rho}(x^p\mathcal{L}^q)) = I$. This means that $\tilde{\rho}(x^p\mathcal{L}^q)$ lies in $\ker\chi = \{(0, 2m\pi) \mid m \in \mathbb{Z}\}$. Hence $\tilde{\rho}(x^p\mathcal{L}^q) = (0, 0)$. Then $\tilde{\rho}$ can induce a homomorphism $\pi_1(K(r)) \to \widetilde{SL_2(\mathbb{R})}$ with non-abelian image. Recall that $\widetilde{SL_2(\mathbb{R})}$ is left-orderable ([2]) and any (non-trivial) subgroup of a left-orderable group is left-orderable. Since $K(r)$ is irreducible [12], $\pi_1(K(r))$ is left-orderable by

[4, Theorem 1.1]. For $r = 0$, $K(0)$ is irreducible and has positive betti number. Hence $\pi_1(K(0))$ is left-orderable by [4, Corollary 3.4]. Thus we have shown that any slope in $[0, 4m)$ is left-orderable for $K = K(m, n)$.

If we apply this argument for $K(n, m)$, then any slope in $[0, 4n)$ is shown to be left-orderable. Since $K(n, m)$ is equivalent to the mirror image of $K(m, n)$, any slope in $(-4n, 0]$ is left-orderable for $K(m, n)$. Thus $I = (-4n, 4m)$ consists of left-orderable slopes for $K = K(m, n)$.

The other cases are proved similarly. □

References

1. V. Bargmann, *Irreducible unitary representations of the Lorentz group*, Ann. of Math. **48** (1947), 568–640.
2. G. Bergman, *Right orderable groups that are not locally indicable*, Pacific J. Math. **147** (1991), 243–248.
3. S. Boyer, C. McA. Gordon and L. Watson, *On L-spaces and left-orderable fundamental groups*, to appear in Math. Ann.
4. S. Boyer, D. Rolfsen and B. Wiest, *Orderable 3-manifold groups*, Ann. Inst. Fourier (Grenoble) **55** (2005), 243–288.
5. G. Burde and H. Zieschang, *Knots*, de Gruyter Studies in Mathematics, 5. Walter de Gruyter & Co., Berlin, 2003.
6. A. Clay, T. Lidman and L. Watson, *Graph manifolds, left-orderability and amalgamation*, preprint, `arXiv:1106.0486`.
7. A. Clay and M. Teragaito, *Left-orderability and exceptional Dehn surgery on two-bridge knots*, to appear in the Proceedings of Geometry and Topology Down Under, Contemporary Mathematics Series.
8. J. Dubois, Y. Huynh and Y. Yamaguchi, *Non-abelian Reidemeister torsion for twist knots*, J. Knot Theory Ramifications **18** (2009), 303–341.
9. E. Ghys, *Groups acting on the circle*, Enseign. Math. **47** (2001), 329–407.
10. R. Hakamata and M. Teragaito, *Left-orderable fundamental group and Dehn surgery on the knot* 5_2, preprint, `arXiv:1208.2087`.
11. R. Hakamata and M. Teragaito, *Left-orderable fundamental group and Dehn surgery on twist knots*, preprint, `arXiv:1212.6305`.
12. A. Hatcher and W. Thurston, *Incompressible surfaces in 2-bridge knot complements*, Invent. Math. **79** (1985), 225–246.
13. J. Hoste and P. Shanahan, *A formula for the A-polynomial of twist knots*, J. Knot Theory Ramifications **13** (2004), 193–209.
14. V. T. Khoi, *A cut-and-paste method for computing the Seifert volumes*, Math. Ann. **326** (2003), 759–801.
15. T. Morifuji and A. T. Tran, *Twisted Alexander polynomilas of* 2*-bridge knots for parabolic representations*, preprint, `arXiv:1301.1101`.
16. P. Ozsváth and Z. Szabó, *On knot Floer homology and lens space surgeries*, Topology **44** (2005), 1281–1300.
17. R. Riley, *Nonabelian representations of 2-bridge knot groups*, Quart. J. Math. Oxford Ser. (2) **35** (1984), 191–208.
18. H. Schubert, *Knoten mit zwei Brücken*, Math. Z. **65** (1956), 133–170.
19. M. Teragaito, *Left-orderability and exceptional Dehn surgery on twist knots*, to appear in Canad. Math. Bull.

20. A. T. Tran, *On left-orderable fundamental groups and Dehn surgeries on twist knots*, preprint.

Graduate School of Education, Hiroshima University, 1-1-1 Kagamiyama, Higashi-hiroshima, Japan 739-8524.

Department of Mathematics and Mathematics Education, Hiroshima University, 1-1-1 Kagamiyama, Higashi-hiroshima, Japan 739-8524.

E-mail address: `teragai@hiroshima-u.ac.jp`